

Below is a list of recent publications that cite LP DAAC data. Please note that this list is not exhaustive or complete.

- Ahmed Suliman, A.H., Gumindoga, W., Katimon, A., and Darus, I.Z.M., 2014, Semi-distributed rainfall-runoff modeling utilizing ASTER DEM in Pinang catchment of Malaysia: *Sains Malaysiana*, v. 43, no. 9, p. 1379-1388. [Also available at http://dx.doi.org/http://www.ukm.my/jsm/pdf_files/SM-PDF-43-9-2014/12%20Ali%20H.%20Ahmad.pdf.]
- Alexander, P.M., Tedesco, M., Fettweis, X., Van De Wal, R.S.W., Smeets, C.J.P.P., and Van Den Broeke, M.R., 2014, Assessing spatio-temporal variability and trends in modelled and measured Greenland Ice Sheet albedo (2000-2013): *Cryosphere*, v. 8, no. 6, p. 2293-2312. [Also available at <http://dx.doi.org/10.5194/tc-8-2293-2014>.]
- Altaf, S., Meraj, G., and Romshoo, S.A., 2014, Morphometry and land cover based multi-criteria analysis for assessing the soil erosion susceptibility of the western Himalayan watershed: *Environmental Monitoring and Assessment*, v. 186, no. 12, p. 8391-8412. [Also available at <http://dx.doi.org/10.1007/s10661-014-4012-2>.]
- Anees, A., and Aryal, J., 2014, A statistical framework for near-real time detection of beetle infestation in pine forests using MODIS data: *IEEE Geoscience and Remote Sensing Letters*, v. 11, no. 10, p. 1717-1721. [Also available at <http://dx.doi.org/10.1109/LGRS.2014.2306712>.]
- Badreldin, N., Frankl, A., and Goossens, R., 2014, Assessing the spatiotemporal dynamics of vegetation cover as an indicator of desertification in Egypt using multi-temporal MODIS satellite images: *Arabian Journal of Geosciences*, v. 7, no. 11, p. 4461-4475. [Also available at <http://dx.doi.org/10.1007/s12517-013-1142-8>.]
- Bhardwaj, A., Joshi, P.K., Snehmani, Singh, M.K., Sam, L., and Gupta, R.D., 2014, Mapping debris-covered glaciers and identifying factors affecting the accuracy: *Cold Regions Science and Technology*, v. 106, p. 161-174. [Also available at <http://dx.doi.org/10.1016/j.coldregions.2014.07.006>.]
- Brando, P.M., Balch, J.K., Nepstad, D.C., Morton, D.C., Putz, F.E., Coe, M.T., Silvério, D., Macedo, M.N., Davidson, E.A., Nóbrega, C.C., Alencar, A., and Soares-Filho, B.S., 2014, Abrupt increases in Amazonian tree mortality due to drought-fire interactions: *Proceedings of the National Academy of Sciences of the United States of America*, v. 111, no. 17, p. 6347-6352. [Also available at <http://dx.doi.org/10.1073/pnas.1305499111>.]
- Broich, M., Huete, A., Tulbure, M.G., Ma, X., Xin, Q., Paget, M., Restrepo-Coupe, N., Davies, K., Devadas, R., and Held, A., 2014, Land surface phenological response to decadal climate variability across Australia using satellite remote sensing: *Biogeosciences*, v. 11, no. 18, p. 5181-5198. [Also available at <http://dx.doi.org/10.5194/bg-11-5181-2014>.]
- Bunting, D.P., Kurc, S.A., Glenn, E.P., Nagler, P.L., and Scott, R.L., 2014, Insights for empirically modeling evapotranspiration influenced by riparian and upland vegetation in semiarid regions: *Journal of Arid Environments*, v. 111, p. 42-52. [Also available at <http://dx.doi.org/10.1016/j.jaridenv.2014.06.007>.]
- Cai, H., Yang, X., Wang, K., and Xiao, L., 2014, Is forest restoration in the southwest China karst promoted mainly by climate change or human-induced factors?: *Remote Sensing*, v. 6, no. 10, p. 9895-9910. [Also available at <http://dx.doi.org/10.3390/Rs6109895>.]
- Campion, R., 2014, New lava lake at Nyamuragira volcano revealed by combined ASTER and OMI SO₂ measurements: *Geophysical Research Letters*, v. 41, no. 21, p. 7485-7492. [Also available at <http://dx.doi.org/10.1002/2014GL061808>.]
- Cardozo, F.S., Pereira, G., Shimabukuro, Y.E., and Moraes, E.C., 2014, Analysis of changes in physical parameters of surface derived from biomass burning in Rondônia state: *Boletim de Ciencias Geodesicas*, v. 20, no. 4, p. 830-845. [Also available at <http://dx.doi.org/10.1590/S1982-21702014000400047>.]
- Chen, L., Jin, Z., Michishita, R., Cai, J., Yue, T., Chen, B., and Xu, B., 2014, Dynamic monitoring of wetland cover changes using time-series remote sensing imagery: *Ecological Informatics*, v. 24, p. 17-26. [Also available at <http://dx.doi.org/10.1016/j.ecoinf.2014.06.007>.]
- Chen, L., Michishita, R., and Xu, B., 2014, Abrupt spatiotemporal land and water changes and their potential drivers in Poyang Lake, 2000-2012: *ISPRS Journal of Photogrammetry and Remote Sensing*, v. 98, p. 85-93. [Also available at <http://dx.doi.org/10.1016/j.isprsjprs.2014.09.014>.]

- Chen, Q., Chen, Y.H., Wang, M.J., Jiang, W.G., Hou, P., and Li, Y., 2014, Change of vegetation net primary productivity in Yellow River watersheds from 2001 to 2010 and its climatic driving factors analysis: Chinese Journal of Applied Ecology, v. 25, no. 10, p. 2811-2818. [Also available at <http://dx.doi.org/http://www.cjae.net/EN/Y2014/V25/I10/2811>.]
- Chen, W., Foy, N., Olthof, I., Zhang, Y., Fraser, R., Latifovic, R., Poitevin, J., Zorn, P., and McLennan, D., 2014, A biophysically based and objective satellite seasonality observation method for applications over the Arctic: International Journal of Remote Sensing, v. 35, no. 18, p. 6742-6763. [Also available at <http://dx.doi.org/10.1080/01431161.2014.963897>.]
- Chen, Y., Wang, B., Pollino, C.A., Cuddy, S.M., Merrin, L.E., and Huang, C., 2014, Estimate of flood inundation and retention on wetlands using remote sensing and GIS: Ecohydrology, v. 7, no. 5, p. 1412-1420. [Also available at <http://dx.doi.org/10.1002/eco.1467>.]
- Colditz, R.R., Pouliot, D., Llamas, R.M., Homer, C., Latifovic, R., Ressl, R.A., Tovar, C.M., Hernandez, A.V., and Richardson, K., 2014, Detection of North American land cover change between 2005 and 2010 with 250m MODIS data: Photogrammetric Engineering and Remote Sensing, v. 80, no. 10, p. 918-924. [Also available at http://dx.doi.org/http://www.asprs.org/a/publications/pers/2014journals/PERS_October_2014/HTML/files/assets/basic-html/toc.html.]
- Corbari, C., and Mancini, M., 2014, Intercomparison across scales between remotely-sensed land surface temperature and representative equilibrium temperature from a distributed energy water balance model: Hydrological Sciences Journal-Journal Des Sciences Hydrologiques, v. 59, no. 10, p. 1830-1843. [Also available at <http://dx.doi.org/10.1080/02626667.2014.946418>.]
- Cui, S., Yang, S., Zhu, C., and Wen, N., 2014, Remote sensing of surface reflective properties—role of regularization and a priori knowledge: Optik, v. 125, no. 23, p. 7106-7112. [Also available at <http://dx.doi.org/10.1016/j.jleo.2014.08.089>.]
- da Silva Junior, C.A., Moreira, E.P., Frank, T., Moreira, M.A., and Barcellos, D., 2014, Comparison of areas of soybean (*Glycine max* (L) Merr.) obtained through the interpretation of TM/Landsat and MODIS/TERRA images in the municipality of Maracaju (MS): Bioscience Journal, v. 30, no. 5, p. 707-716. [Also available at <http://dx.doi.org/http://www.seer.ufu.br/index.php/biosciencejournal/article/view/19738>.]
- Danelichen, V.H.M., Biudes, M.S., Souza, M.C., Machado, N.G., Da Silva, B.B., and Nogueira, J.S., 2014, Estimation of soil heat flux in a neotropical wetland region using remote sensing techniques: Revista Brasileira de Meteorologia, v. 29, no. 4, p. 469-482. [Also available at <http://dx.doi.org/10.1590/0102-778620120568>.]
- Dempewolf, J., Adusei, B., Becker-Reshef, I., Hansen, M., Potapov, P., Khan, A., and Barker, B., 2014, Wheat Yield Forecasting for Punjab Province from Vegetation Index Time Series and Historic Crop Statistics: Remote Sensing, v. 6, no. 10, p. 9653-9675. [Also available at <http://dx.doi.org/10.3390/rs6109653>.]
- Dente, L., Ferrazzoli, P., Su, Z., van der Velde, R., and Guerriero, L., 2014, Combined use of active and passive microwave satellite data to constrain a discrete scattering model: Remote Sensing of Environment, v. 155, p. 222-238. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.031>.]
- Di Mauro, B., Fava, F., Busetto, L., Crosta, G.F., and Colombo, R., 2014, Post-fire resilience in the Alpine region estimated from MODIS satellite multispectral data: International Journal of Applied Earth Observation and Geoinformation, v. 32, p. 163-172. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.04.010>.]
- Ding, Y., Ge, Y., Hu, M., Wang, J., Wang, J., Zheng, X., and Zhao, K., 2014, Comparison of spatial sampling strategies for ground sampling and validation of MODIS LAI products: International Journal of Remote Sensing, v. 35, no. 20, p. 7230-7244. [Also available at <http://dx.doi.org/10.1080/01431161.2014.967889>.]
- Elsaid, M., Aboelkhair, H., Dardier, A., and Hermas, E., 2014, Investigation of a relation between radiogenic heat production and kinetic surface temperature from multispectral ASTER-TIR data—a case study on Elmissikat-Eleridiya granites, Central Eastern Desert, Egypt: Arabian Journal of Geosciences, v. 7, no. 11, p. 4615-4628. [Also available at <http://dx.doi.org/10.1007/s12517-013-1118-8>.]

- Emel, J., Plisinski, J., and Rogan, J., 2014, Monitoring geomorphic and hydrologic change at mine sites using satellite imagery—the Geita Gold Mine in Tanzania: *Applied Geography*, v. 54, p. 243-249. [Also available at <http://dx.doi.org/10.1016/j.apgeog.2014.07.009>.]
- Ermolaev, O.P., Mal'tsev, K.A., and Ivanov, M.A., 2014, Automated construction of the boundaries of basin geosystems for the Volga Federal District: *Geography and Natural Resources*, v. 35, no. 3, p. 222-228. [Also available at <http://dx.doi.org/10.1134/S1875372814030044>.]
- Estrada-Peña, A., Tarragona, E.L., Vesco, U., Meneghi, D.D., Mastropaoletti, M., Mangold, A.J., Guglielmone, A.A., and Nava, S., 2014, Divergent environmental preferences and areas of sympatry of tick species in the Amblyomma cajennense complex (Ixodidae): *International Journal for Parasitology*, v. 44, no. 14, p. 1081-1089. [Also available at <http://dx.doi.org/10.1016/j.ijpara.2014.08.007>.]
- Fan, X.M., Liu, H.G., Liu, G.H., and Li, S.B., 2014, Reconstruction of MODIS land-surface temperature in a flat terrain and fragmented landscape: *International Journal of Remote Sensing*, v. 35, no. 23, p. 7857-7877. [Also available at <http://dx.doi.org/10.1080/01431161.2014.978036>.]
- Fang, S., Le, Y., Liang, Q., and Liu, X., 2014, Leaf area index estimation using time-series MODIS Data in different types of vegetation: *Journal of the Indian Society of Remote Sensing*, v. 42, no. 4, p. 733-743. [Also available at <http://dx.doi.org/10.1007/s12524-013-0349-1>.]
- Faruolo, M., Coviello, I., Filizzola, C., Lacava, T., Pergola, N., and Tramutoli, V., 2014, A satellite-based analysis of the Val d'Agri Oil Center (southern Italy) gas flaring emissions: *Natural Hazards and Earth System Sciences*, v. 14, no. 10, p. 2783-2793. [Also available at <http://dx.doi.org/10.5194/nhess-14-2783-2014>.]
- Feidas, H., Karagiannidis, A., Keppas, S., Vaitis, M., Kontos, T., Zanis, P., Melas, D., and Anadranistakis, E., 2014, Modeling and mapping temperature and precipitation climate data in Greece using topographical and geographical parameters: *Theoretical and Applied Climatology*, v. 118, no. 1-2, p. 133-146. [Also available at <http://dx.doi.org/10.1007/s00704-013-1052-4>.]
- Franch, B., Vermote, E.F., Sobrino, J.A., and Julien, Y., 2014, Retrieval of surface albedo on a daily basis—application to MODIS data: *IEEE Transactions on Geoscience and Remote Sensing*, v. 52, no. 12, p. 7549-7558. [Also available at <http://dx.doi.org/10.1109/TGRS.2014.2313842>.]
- French, N.H.F., McKenzie, D., Erickson, T., Koziol, B., Billmire, M., Arthur Endsley, K., Yager Scheinerman, N.K., Jenkins, L., Miller, M.E., Ottmar, R., and Prichard, S., 2014, Modeling regional-scale wildland fire emissions with the Wildland Fire Emissions Information System: *Earth Interactions*, v. 18, no. 16. [Also available at <http://dx.doi.org/10.1175/EI-D-14-0002.1>.]
- Fu, Q., Min, X., Sun, L., and Ma, S., 2014, Atmospheric correction of HJ-1 A/B CCD over land—land surface reflectance calculation for geographical information product: *Journal of Geographical Sciences*, v. 24, no. 6, p. 1083-1094. [Also available at <http://dx.doi.org/10.1007/s11442-014-1140-0>.]
- Fu, Y., Zhang, H., Dong, W., and Yuan, W., 2014, Comparison of phenology models for predicting the onset of growing season over the northern hemisphere: *PLoS ONE*, v. 9, no. 10. [Also available at <http://dx.doi.org/10.1371/journal.pone.0109544>.]
- Fu, Y.H., Piao, S., Zhao, H., Jeong, S.-J., Wang, X., Vitasse, Y., Ciais, P., and Janssens, I.A., 2014, Unexpected role of winter precipitation in determining heat requirement for spring vegetation green-up at northern middle and high latitudes: *Global Change Biology*, v. 20, no. 12, p. 3743-3755. [Also available at <http://dx.doi.org/10.1111/gcb.12610>.]
- Fuchs, M., Torizin, J., and Kühn, F., 2014, The effect of DEM resolution on the computation of the factor of safety using an infinite slope model: *Geomorphology*, v. 224, p. 16-26. [Also available at <http://dx.doi.org/10.1016/j.geomorph.2014.07.015>.]
- Fuentes, S., Poblete-Echeverria, C., Ortega-Farias, S., Tyerman, S., and De Bei, R., 2014, Automated estimation of leaf area index from grapevine canopies using cover photography, video and computational analysis methods: *Australian Journal of Grape and Wine Research*, v. 20, no. 3, p. 465-473. [Also available at <http://dx.doi.org/10.1111/ajgw.12098>.]

- Gamarra, N.L.R., Corrêa, M.P., and Targino, A.C.L., 2014, Use of remote sensing to retrieve surface albedo and land surface temperature in Londrina (Paraná)—a contribution to urban heat island studies: Revista Brasileira de Meteorologia, v. 29, no. 4, p. 537-550. [Also available at <http://dx.doi.org/10.1590/0102-778620130671>.]
- Gan, W., Shen, H., Zhang, L., and Gong, W., 2014, Normalization of medium-resolution NDVI by the use of coarser reference data—method and evaluation: International Journal of Remote Sensing, v. 35, no. 21, p. 7400-7429. [Also available at <http://dx.doi.org/10.1080/01431161.2014.968684>.]
- Gao, S., Zhu, Z., Liu, S., Jin, R., Yang, G., and Tan, L., 2014, Estimating the spatial distribution of soil moisture based on Bayesian maximum entropy method with auxiliary data from remote sensing: International Journal of Applied Earth Observation and Geoinformation, v. 32, no. 1, p. 54-66. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.03.003>.]
- Gideon Neba, S., Kanninen, M., Eba'a Atyi, R., and Sonwa, D.J., 2014, Assessment and prediction of above-ground biomass in selectively logged forest concessions using field measurements and remote sensing data—case study in South East Cameroon: Forest Ecology and Management, v. 329, p. 177-185. [Also available at <http://dx.doi.org/10.1016/j.foreco.2014.06.018>.]
- Giri, C., and Long, J., 2014, Land cover characterization and mapping of South America for the year 2010 using Landsat 30m satellite data: Remote Sensing, v. 6, no. 10, p. 9494-9510. [Also available at <http://dx.doi.org/10.3390/rs6109494>.]
- Glanz, H., Carvalho, L., Sulla-Menashe, D., and Friedl, M.A., 2014, A parametric model for classifying land cover and evaluating training data based on multi-temporal remote sensing data: ISPRS Journal of Photogrammetry and Remote Sensing, v. 97, p. 219-228. [Also available at <http://dx.doi.org/10.1016/j.isprsjprs.2014.09.004>.]
- Gorchakov, G.I., Sitnov, S.A., Sviridenkov, M.A., Semoutnikova, E.G., Emilenko, A.S., Isakov, A.A., Kopeikin, V.M., Karpov, A.V., Gorchakova, I.A., Verichev, K.S., Kurbatov, G.A., and Ponomareva, T.Y., 2014, Satellite and ground-based monitoring of smoke in the atmosphere during the summer wildfires in European Russia in 2010 and Siberia in 2012: International Journal of Remote Sensing, v. 35, no. 15, p. 5698-5721. [Also available at <http://dx.doi.org/10.1080/01431161.2014.945008>.]
- Gou, Y., Wu, W., Liu, H., Guo, P., Wang, X., and Xie, D., 2014, Rice production estimation using empirical modeling with MODIS NDVI data in Southwestern China: ICIC Express Letters, Part B—Applications, v. 5, no. 6, p. 1727-1733. [Also available at <http://dx.doi.org/http://www.ijicic.org/elb-5%286%29.htm>.]
- Guay, K.C., Beck, P.S.A., Berner, L.T., Goetz, S.J., Baccini, A., and Buermann, W., 2014, Vegetation productivity patterns at high northern latitudes—a multi-sensor satellite data assessment: Global Change Biology, v. 20, no. 10, p. 3147-3158. [Also available at <http://dx.doi.org/10.1111/gcb.12647>.]
- Guillevic, P.C., Biard, J.C., Hulley, G.C., Privette, J.L., Hook, S.J., Olioso, A., Goettsche, F.M., Radocinski, R., Roman, M.O., Yu, Y., and Csizsar, I., 2014, Validation of Land Surface Temperature products derived from the Visible Infrared Imaging Radiometer Suite (VIIRS) using ground-based and heritage satellite measurements: Remote Sensing of Environment, v. 154, p. 19-37. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.013>.]
- Guillevic, P.C., Biard, J.C., Hulley, G.C., Privette, J.L., Hook, S.J., Olioso, A., Götsche, F.M., Radocinski, R., Román, M.O., Yu, Y., and Csizsar, I., 2014, Validation of Land Surface Temperature products derived from the Visible Infrared Imaging Radiometer Suite (VIIRS) using ground-based and heritage satellite measurements: Remote Sensing of Environment, v. 154, p. 19-37. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.013>.]
- Guindon, L., Bernier, P.Y., Beaudoin, A., Pouliot, D., Villemaire, P., Hall, R.J., Latifovic, R., and St-Amant, R., 2014, Annual mapping of large forest disturbances across Canada's forests using 250 m MODIS imagery from 2000 to 2011: Canadian Journal of Forest Research, v. 44, no. 12, p. 1545-1554. [Also available at <http://dx.doi.org/10.1139/cjfr-2014-0229>.]
- Guzinski, R., Nieto, H., Jensen, R., and Mendiguren, G., 2014, Remotely sensed land-surface energy fluxes at sub-field scale in heterogeneous agricultural landscape and coniferous plantation: Biogeosciences, v. 11, no. 18, p. 5021-5046. [Also available at <http://dx.doi.org/10.5194/bg-11-5021-2014>.]
- Hammer, D., Kraft, R., and Wheeler, D., 2014, Alerts of forest disturbance from MODIS imagery: International Journal of Applied Earth Observation and Geoinformation, v. 33, p. 1-9. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.04.011>.]

He, T., Liang, S., and Song, D.-X., 2014, Analysis of global land surface albedo climatology and spatial-temporal variation during 1981–2010 from multiple satellite products: *Journal of Geophysical Research-Atmospheres*, v. 119, no. 17. [Also available at <http://dx.doi.org/10.1002/2014jd021667>.]

Heiblum, R.H., Koren, I., and Feingold, G., 2014, On the link between Amazonian forest properties and shallow cumulus cloud fields: *Atmospheric Chemistry and Physics*, v. 14, no. 12, p. 6063-6074. [Also available at <http://dx.doi.org/10.5194/acp-14-6063-2014>.]

Hereher, M.E., 2014, Capacity assessment of the Qattara Depression—Egypt as a sink for the global sea level rise: *Geocarto International*. [Also available at <http://dx.doi.org/10.1080/10106049.2014.966159>.]

Herron-Thorpe, F.L., Mount, G.H., Emmons, L.K., Lamb, B.K., Jaffe, D.A., Wigder, N.L., Chung, S.H., Zhang, R., Woelfle, M.D., and Vaughan, J.K., 2014, Air quality simulations of wildfires in the Pacific Northwest evaluated with surface and satellite observations during the summers of 2007 and 2008: *Atmospheric Chemistry and Physics*, v. 14, no. 22, p. 12533-12551. [Also available at <http://dx.doi.org/10.5194/acp-14-12533-2014>.]

Hilker, T., Lyapustin, A.I., Tucker, C.J., Hall, F.G., Myneni, R.B., Wang, Y., Bi, J., De Moura, Y.M., and Sellers, P.J., 2014, Vegetation dynamics and rainfall sensitivity of the Amazon: *Proceedings of the National Academy of Sciences of the United States of America*, v. 111, no. 45, p. 16041-16046. [Also available at <http://dx.doi.org/10.1073/pnas.1404870111>.]

Holzman, M.E., Rivas, R., and Bayala, M., 2014, Subsurface soil moisture estimation by VI-LST method: *IEEE Geoscience and Remote Sensing Letters*, v. 11, no. 11, p. 1951-1955. [Also available at <http://dx.doi.org/10.1109/lgrs.2014.2314617>.]

Hu, S., Mo, X., and Lin, Z., 2014, Optimizing the photosynthetic parameter Vcmax by assimilating MODIS-fPAR and MODIS-NDVI with a process-based ecosystem model: *Agricultural and Forest Meteorology*, v. 198, p. 320-334. [Also available at <http://dx.doi.org/10.1016/j.agrformet.2014.09.002>.]

Huang, B., and Zhang, H., 2014, Spatio-temporal reflectance fusion via unmixing—accounting for both phenological and land-cover changes: *International Journal of Remote Sensing*, v. 35, no. 16, p. 6213-6233. [Also available at <http://dx.doi.org/10.1080/01431161.2014.951097>.]

Huang, N., Gu, L., and Niu, Z., 2014, Estimating soil respiration using spatial data products—a case study in a deciduous broadleaf forest in the Midwest USA: *Journal of Geophysical Research-Atmospheres*, v. 119, no. 11, p. 6393-6408. [Also available at <http://dx.doi.org/10.1002/2013JD020515>.]

Hui, S., Mo, X., and Lin, Z., 2014, Optimizing the photosynthetic parameter V-cmax by assimilating MODIS-f(PAR) and MODIS-NDVI with a process-based ecosystem model: *Agricultural and Forest Meteorology*, v. 198, p. 320-334. [Also available at <http://dx.doi.org/10.1016/j.agrformet.2014.09.002>.]

Jacques, D.C., Kergoat, L., Hiernaux, P., Mougin, E., and Defourny, P., 2014, Monitoring dry vegetation masses in semi-arid areas with MODIS SWIR bands: *Remote Sensing of Environment*, v. 153, p. 40-49. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.07.027>.]

Ji, L., Zhang, L., Rover, J., Wylie, B.K., and Chen, X., 2014, Geostatistical estimation of signal-to-noise ratios for spectral vegetation indices: *ISPRS Journal of Photogrammetry and Remote Sensing*, v. 96, p. 20-27. [Also available at <http://dx.doi.org/10.1016/j.isprsjprs.2014.06.013>.]

Jia, K., Liang, S., Wei, X., Yao, Y., Su, Y., Jiang, B., and Wang, X., 2014, Land cover classification of Landsat data with phenological features extracted from time series MODIS NDVI data: *Remote Sensing*, v. 6, no. 11, p. 11518-11532. [Also available at <http://dx.doi.org/10.3390/rs61111518>.]

Jia, K., Liang, S., Zhang, L., Wei, X., Yao, Y., and Xie, X., 2014, Forest cover classification using Landsat ETM plus data and time series MODIS NDVI data: *International Journal of Applied Earth Observation and Geoinformation*, v. 33, p. 32-38. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.04.015>.]

- Jia, K., Liang, S., Zhang, L., Wei, X., Yao, Y., and Xie, X., 2014, Forest cover classification using Landsat ETM+ data and time series MODIS NDVI data: International Journal of Applied Earth Observation and Geoinformation, v. 33, p. 32-38. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.04.015>.]
- Jiang, G.M., and Liu, R., 2014, Retrieval of sea and land surface temperature from SVISSR/FY-2C/D/E measurements: IEEE Transactions on Geoscience and Remote Sensing, v. 52, no. 10, p. 6132-6140. [Also available at <http://dx.doi.org/10.1109/TGRS.2013.2295260>.]
- Jiang, J., Xiao, Z., Wang, J., and Song, J., 2014, Sequential method with incremental analysis update to retrieve leaf area index from time series MODIS reflectance data: Remote Sensing, v. 6, no. 10, p. 9194-9212. [Also available at <http://dx.doi.org/10.3390/Rs6109194>.]
- Jin, X.M., Guo, R.H., Zhang, Q., Zhou, Y.X., Zhang, D.R., and Yang, Z., 2014, Response of vegetation pattern to different landform and water-table depth in Hailiutu River basin, Northwestern China: Environmental Earth Sciences, v. 71, no. 11, p. 4889-4898. [Also available at <http://dx.doi.org/10.1007/s12665-013-2882-1>.]
- Jing, C., Shortridge, A., Lin, S., and Wu, J., 2014, Comparison and validation of SRTM and ASTER GDEM for a subtropical landscape in Southeastern China: International Journal of Digital Earth, v. 7, no. 12, p. 969-992. [Also available at <http://dx.doi.org/10.1080/17538947.2013.807307>.]
- Karnieli, A., Qin, Z., Wu, B., Panov, N., and Yan, F., 2014, Spatio-temporal dynamics of land-use and land-cover in the Mu Us Sandy Land, China, using the change vector analysis technique: Remote Sensing, v. 6, no. 10, p. 9316-9339. [Also available at <http://dx.doi.org/10.3390/Rs6109316>.]
- Ke, L., and Song, C., 2014, Remotely sensed surface temperature variation of an inland saline lake over the central qinghai-tibet plateau: ISPRS Journal of Photogrammetry and Remote Sensing, v. 98, p. 157-167. [Also available at <http://dx.doi.org/10.1016/j.isprsjprs.2014.09.007>.]
- Kellogg, C.H., and Zhou, X., 2014, Impact of the construction of a large dam on riparian vegetation cover at different elevation zones as observed from remotely sensed data: International Journal of Applied Earth Observation and Geoinformation, v. 32, p. 19-34. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.03.021>.]
- Kim, J., Grunwald, S., and Rivero, R.G., 2014, Soil phosphorus and nitrogen predictions across spatial escalating scales in an aquatic ecosystem using remote sensing images: IEEE Transactions on Geoscience and Remote Sensing, v. 52, no. 10, p. 6724-6737. [Also available at <http://dx.doi.org/10.1109/tgrs.2014.2301443>.]
- Klisch, A., and Atzberger, C., 2014, Evaluating phenological metrics derived from the MODIS time series over the European continent: Photogrammetrie, Fernerkundung, Geoinformation, v. 2014, no. 5, p. 409-421. [Also available at <http://dx.doi.org/10.1127/1432-8364/2014/0233>.]
- Koerner, P., Pluntke, T., Sachse, A., Boettcher, N., Naumov, D., Kolditz, O., and Bernhofer, C., 2014, Inverse determination of groundwater inflow using water balance simulations: Environmental Earth Sciences, v. 72, no. 12, p. 4757-4769. [Also available at <http://dx.doi.org/10.1007/s12665-014-3327-1>.]
- Konovalov, I.B., Berezin, E.V., Ciais, P., Broquet, G., Beekmann, M., Hadji-Lazaro, J., Clerbaux, C., Andreae, M.O., Kaiser, J.W., and Schulze, E.D., 2014, Constraining CO₂emissions from open biomass burning by satellite observations of co-emitted species—a method and its application to wildfires in Siberia: Atmospheric Chemistry and Physics, v. 14, no. 19, p. 10383-10410. [Also available at <http://dx.doi.org/10.5194/acp-14-10383-2014>.]
- Kouadio, L., Newlands, N.K., Davidson, A., Zhang, Y., and Chipanshi, A., 2014, Assessing the Performance of MODIS NDVI and EVI for Seasonal Crop Yield Forecasting at the Ecodistrict Scale: Remote Sensing, v. 6, no. 10, p. 10193-10214. [Also available at <http://dx.doi.org/10.3390/Rs61010193>.]
- Krylov, A., McCarty, J.L., Potapov, P., Loboda, T., Tyukavina, A., Turubanova, S., and Hansen, M.C., 2014, Remote sensing estimates of stand-replacement fires in Russia, 2002-2011: Environmental Research Letters, v. 9, no. 10. [Also available at <http://dx.doi.org/10.1088/1748-9326/9/10/105007>.]

Kumar, A.S., Radhika, T., Saritha, P.K., Keerthi, V., Anjani, R.N., Kumar, M.S., Sekhar, K.S., Satyanarayana, P., Sudha, M.S.N., Sai, M.V.R.S., and Dadhwal, V.K., 2014, Generation of vegetation fraction and surface albedo products over India from Ocean Colour Monitor (OCM) data onboard Oceansat-2: Journal of the Indian Society of Remote Sensing, v. 42, no. 4, p. 701-709. [Also available at <http://dx.doi.org/10.1007/s12524-014-0371-y>.]

La Sorte, F.A., Fink, D., Hochachka, W.M., DeLong, J.P., and Kelling, S., 2014, Spring phenology of ecological productivity contributes to the use of looped migration strategies by birds: Proceedings of the Royal Society B-Biological Sciences, v. 281, no. 1793. [Also available at <http://dx.doi.org/10.1098/rspb.2014.0984>.]

Landmann, T., and Dubovyk, O., 2014, Spatial analysis of human-induced vegetation productivity decline over eastern Africa using a decade (2001-2011) of medium resolution MODIS time-series data: International Journal of Applied Earth Observation and Geoinformation, v. 33, p. 76-82. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.04.020>.]

Laxmi, K., and Nandagiri, L., 2014, Latent heat flux estimation using trapezoidal relationship between MODIS land surface temperature and fraction of vegetation-application and validation in a humid tropical region: Remote Sensing Letters, v. 5, no. 11, p. 981-990. [Also available at <http://dx.doi.org/10.1080/2150704X.2014.984083>.]

Lazzarini, M., Marpu, P.R., Eissa, Y., and Ghedira, H., 2014, Toward a near real-time product of air temperature maps from satellite data and in situ measurements in arid environments: IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, v. 7, no. 7, p. 3093-3104. [Also available at <http://dx.doi.org/10.1109/JSTARS.2014.2320762>.]

Le, T.H., Thanh Nguyen, T.N., Lasko, K., Ilavajhala, S., Vadrevu, K.P., and Justice, C., 2014, Vegetation fires and air pollution in Vietnam: Environmental Pollution, v. 195, p. 267-275. [Also available at <http://dx.doi.org/10.1016/j.envpol.2014.07.023>.]

Letu, H., Nagao, T.M., Nakajima, T.Y., and Matsumae, Y., 2014, Method for validating cloud mask obtained from satellite measurements using ground-based sky camera: Applied Optics, v. 53, no. 31, p. 7523-7533. [Also available at <http://dx.doi.org/10.1364/ao.53.007523>.]

Li, C.-f., Shen, D., Dong, J.-s., Yin, J.-y., Zhao, J.-j., and Xue, D., 2014, Monitoring of urban heat island in Shanghai, China, from 1981 to 2010 with satellite data: Arabian Journal of Geosciences, v. 7, no. 10, p. 3961-3971. [Also available at <http://dx.doi.org/10.1007/s12517-013-1053-8>.]

Li, G.Y., Li, X.Y., Zhao, G.Q., Zhang, Z.H., and Li, Y.T., 2014, Characteristics of spatial and temporal phenology under the dynamic variation of grassland in the Qinghai Lake watershed: Shengtai Xuebao/ Acta Ecologica Sinica, v. 34, no. 11. [Also available at <http://dx.doi.org/10.5846/stxb201211251668>.]

Li, M., Song, Y., Huang, X., Li, J., Mao, Y., Zhu, T., Cai, X., and Liu, B., 2014, Improving mesoscale modeling using satellite-derived land surface parameters in the Pearl River Delta region, China: Journal of Geophysical Research-Atmospheres, v. 119, no. 11, p. 6325-6346. [Also available at <http://dx.doi.org/10.1002/2014JD021871>.]

Li, Q., Cao, X., Jia, K., Zhang, M., and Dong, Q., 2014, Crop type identification by integration of high-spatial resolution multispectral data with features extracted from coarse-resolution time-series vegetation index data: International Journal of Remote Sensing, v. 35, no. 16, p. 6076-6088. [Also available at <http://dx.doi.org/10.1080/01431161.2014.943325>.]

Li, X., Wang, J., Song, W., Ma, J., Telesca, L., and Zhang, Y., 2014, Automatic smoke detection in MODIS satellite data based on K-means clustering and fisher linear discrimination: Photogrammetric Engineering and Remote Sensing, v. 80, no. 10, p. 971-982. [Also available at <http://dx.doi.org/10.14358/PERS.80.10.971>.]

Li, Y., Pan, X., Wang, C., Liu, Y., and Zhao, Q., 2014, Changes of vegetation net primary productivity and its driving factors from 2000 to 2011 in Guangxi, China: Shengtai Xuebao/ Acta Ecologica Sinica, v. 34, no. 18, p. 5220-5228. [Also available at <http://dx.doi.org/10.5846/stxb201405100952>.]

---, 2014, Monitoring changes of soil organic matter and total nitrogen in cultivated land in Guangxi by remote sensing: Shengtai Xuebao/ Acta Ecologica Sinica, v. 34, no. 18, p. 5283-5291. [Also available at <http://dx.doi.org/10.5846/stxb201405100953>.]

Liang, L., Schwartz, M.D., Wang, Z., Gao, F., Schaaf, C.B., Tan, B., Morisette, J.T., and Zhang, X., 2014, A Cross comparison of spatiotemporally enhanced springtime phenological measurements from satellites and ground in a northern U.S. mixed forest: *IEEE Transactions on Geoscience and Remote Sensing*, v. 52, no. 12, p. 7513-7526. [Also available at <http://dx.doi.org/10.1109/TGRS.2014.2313558>.]

Liao, Q.F., Zhang, X., Ma, Q., Yao, Y., and Yu, D.P., 2014, Spatiotemporal variation of fractional vegetation cover and remote sensing monitoring in the eastern agricultural region of Qinghai Province: *Shengtai Xuebao/ Acta Ecologica Sinica*, v. 34, no. 20, p. 5936-5943. [Also available at <http://dx.doi.org/10.5846/stxb201301230138>.]

Lin, J., Liu, X., Li, K., and Li, X., 2014, A maximum entropy method to extract urban land by combining MODIS reflectance, MODIS NDVI, and DMSP-OLS data: *International Journal of Remote Sensing*, v. 35, no. 18, p. 6708-6727. [Also available at <http://dx.doi.org/10.1080/01431161.2014.960623>.]

Liu, L.W., Zhang, W.P., Duan, Y.H., Xing, L.X., Chen, S.B., and Zhao, M., 2014, Terrain corrected TVDI for agricultural drought monitoring using MODIS data: *Shengtai Xuebao/ Acta Ecologica Sinica*, v. 34, no. 13, p. 3704-3711. [Also available at <http://dx.doi.org/10.5846/stxb201211221652>.]

Liu, Y., Wang, J., Zhou, H., and Xue, H., 2014, Upscaling approach for validation of LAI products derived from remote sensing observation: *Journal of Remote Sensing*, v. 18, no. 6, p. 1189-1198. [Also available at <http://dx.doi.org/10.11834/jrs.20144084>.]

Liu, Z., Shao, Q., Tao, J., and Chi, W., 2014, Intra-annual variability of satellite observed surface albedo associated with typical land cover types in China: *Journal of Geographical Sciences*, v. 25, no. 1, p. 35-44. [Also available at <http://dx.doi.org/10.1007/s11442-015-1151-5>.]

Liu, Z., Wang, L., and Wang, S., 2014, Comparison of different GPP models in China using MODIS image and ChinaFLUX data: *Remote Sensing*, v. 6, no. 10, p. 10215-10231. [Also available at <http://dx.doi.org/10.3390/Rs61010215>.]

Lu, L., Luo, G.P., and Wang, J.Y., 2014, Development of an ATI-NDVI method for estimation of soil moisture from MODIS data: *International Journal of Remote Sensing*, v. 35, no. 10, p. 3797-3815. [Also available at <http://dx.doi.org/10.1080/01431161.2014.919677>.]

Lyapustin, A., Wang, Y., Xiong, X., Meister, G., Platnick, S., Levy, R., Franz, B., Korkin, S., Hilker, T., Tucker, J., Hall, F., Sellers, P., Wu, A., and Angal, A., 2014, Scientific impact of MODIS C5 calibration degradation and C6+ improvements: *Atmospheric Measurement Techniques*, v. 7, no. 12, p. 4353-4365. [Also available at <http://dx.doi.org/10.5194/amt-7-4353-2014>.]

Ma, X., Huete, A., Yu, Q., Restrepo-Coupe, N., Beringer, J., Hutley, L.B., Kanniah, K.D., Cleverly, J., and Eamus, D., 2014, Parameterization of an ecosystem light-use-efficiency model for predicting savanna GPP using MODIS EVI: *Remote Sensing of Environment*, v. 154, p. 253-271. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.025>.]

Maeda, E.E., Heiskanen, J., Aragão, L.E.O.C., and Rinne, J., 2014, Can MODIS EVI monitor ecosystem productivity in the Amazon rainforest?: *Geophysical Research Letters*, v. 41, no. 20, p. 7176-7183. [Also available at <http://dx.doi.org/10.1002/2014GL061535>.]

Mao, D., Wang, Z., Wu, C., Song, K., and Ren, C., 2014, Examining forest net primary productivity dynamics and driving forces in northeastern China during 1982–2010: *Chinese Geographical Science*, v. 24, no. 6, p. 631-646. [Also available at <http://dx.doi.org/10.1007/s11769-014-0662-9>.]

Martínez, B., Albargues, E., Camacho, F., Moreno, A., and Gilabert, M.A., 2014, fAPAR estimates over the Iberian Peninsula by the inversion of the 4SAIL 2 radiative transfer model: *Revista de Teledetección*, no. 42, p. 61-77. [Also available at <http://dx.doi.org/10.4995/raet.2014.3177>.]

Melkonian, A.K., Willis, M.J., and Pritchard, M.E., 2014, Satellite-derived volume loss rates and glacier speeds for the Juneau Icefield, Alaska: *Journal of Glaciology*, v. 60, no. 222, p. 743-760. [Also available at <http://dx.doi.org/10.3189/2014jog13j181>.]

Micheletty, P.D., Kinoshita, A.M., and Hogue, T.S., 2014, Application of MODIS snow cover products—wildfire impacts on snow and melt in the Sierra Nevada: *Hydrology and Earth System Sciences*, v. 18, no. 11, p. 4601-4615. [Also available at <http://dx.doi.org/10.5194/hess-18-4601-2014.1>]

Mielke, C., Boesche, N.K., Rogass, C., Kaufmann, H., Gauert, C., and de Wit, M., 2014, Spaceborne mine waste mineralogy monitoring in South Africa, applications for modern push-broom missions—Hyperion/OLI and EnMAP/Sentinel-2: *Remote Sensing*, v. 6, no. 8, p. 6790-6816. [Also available at <http://dx.doi.org/10.3390/rs6086790.1>]

Miernecki, M., Wigneron, J.P., Lopez-Baeza, E., Kerr, Y., De Jeu, R., De Lannoy, G.J.M., Jackson, T.J., O'Neill, P.E., Schwank, M., Moran, R.F., Bircher, S., Lawrence, H., Mialon, A., Al Bitar, A., and Richaume, P., 2014, Comparison of SMOS and SMAP soil moisture retrieval approaches using tower-based radiometer data over a vineyard field: *Remote Sensing of Environment*, v. 154, no. 1, p. 89-101. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.002.1>]

Mishra, A.K., 2014, Retrieval of EVI from Oceansat 2 data and comparison with MODIS derived EVI: *Journal of the Indian Society of Remote Sensing*, v. 42, no. 4, p. 877-883. [Also available at <http://dx.doi.org/10.1007/s12524-014-0369-5.1>]

Mitchell, B.C., and Chakraborty, J., Urban heat and climate justice—a landscape of thermal inequity in Pinellas County, Florida: *Geographical Review*, v. 104, no. 4, p. 459-480. [Also available at <http://dx.doi.org/10.1111/j.1931-0846.2014.12039.x.1>]

Monaghan, A.J., Hu, L., Brunsell, N.A., Barlage, M., and Wilhelmi, O.V., 2014, Evaluating the impact of urban morphology configurations on the accuracy of urban canopy model temperature simulations with MODIS: *Journal of Geophysical Research-Atmospheres*, v. 119, no. 11, p. 6376-6392. [Also available at <http://dx.doi.org/10.1002/2013JD021227.1>]

Nagai, S., Ishii, R., Bin Suhaili, A., Kobayashi, H., Matsuoka, M., Ichie, T., Motohka, T., Kendawang, J.J., and Suzuki, R., 2014, Usability of noise-free daily satellite-observed green-red vegetation index values for monitoring ecosystem changes in Borneo: *International Journal of Remote Sensing*, v. 35, no. 23, p. 7910-7926. [Also available at <http://dx.doi.org/10.1080/01431161.2014.978039.1>]

Ni, W., Zhang, Z., Sun, G., Guo, Z., and He, Y., 2014, The penetration depth derived from the synthesis of ALOS/PALSAR InSAR data and ASTER GDEM for the mapping of forest biomass: *Remote Sensing*, v. 6, no. 8, p. 7303-7319. [Also available at <http://dx.doi.org/10.3390/rs6087303.1>]

Nischitha, V., Ahmed, S.A., Varikoden, H., and Revadekar, J.V., 2014, The impact of seasonal rainfall variability on NDVI in the Tunga and Bhadra river basins, Karnataka, India: *International Journal of Remote Sensing*, v. 35, no. 23, p. 8025-8043. [Also available at <http://dx.doi.org/10.1080/01431161.2014.979301.1>]

Noguchi, K., Richter, A., Rozanov, V., Rozanov, A., Burrows, J.P., Irie, H., and Kita, K., 2014, Effect of surface BRDF of various land cover types on geostationary observations of tropospheric NO₂: *Atmospheric Measurement Techniques*, v. 7, no. 10, p. 3497-3508. [Also available at <http://dx.doi.org/10.5194/amt-7-3497-2014.1>]

Othman, A.A., and Gloaguen, R., 2014, Improving lithological mapping by SVM classification of spectral and morphological features—the discovery of a new chromite body in the Mawat Ophiolite Complex (Kurdistan, NE Iraq): *Remote Sensing*, v. 6, no. 8, p. 6867-6896. [Also available at <http://dx.doi.org/10.3390/rs6086867.1>]

Painemal, D., Kato, S., and Minnis, P., 2014, Boundary layer regulation in the southeast Atlantic cloud microphysics during the biomass burning season as seen by the A-train satellite constellation: *Journal of Geophysical Research-Atmospheres*, v. 119, no. 19, p. 11288-11302. [Also available at <http://dx.doi.org/10.1002/2014jd022182.1>]

Pandey, A.K., Singh, S., Berwal, S., Kumar, D., Pandey, P., Prakash, A., Lodhi, N., Maithani, S., Jain, V.K., and Kumar, K., 2014, Spati - temporal variations of urban heat island over Delhi: *Urban Climate*, v. 10, Part 1, p. 119-133. [Also available at <http://dx.doi.org/10.1016/j.uclim.2014.10.005.1>]

Pena-Arancibia, J.L., McVicar, T.R., Paydar, Z., Li, L., Guerschman, J.P., Donohue, R.J., Dutta, D., Podger, G.M., van Dijk, A.I.J.M., and Chiew, F.H.S., 2014, Dynamic identification of summer cropping irrigated areas in a large basin experiencing extreme climatic variability: *Remote Sensing of Environment*, v. 154, p. 139-152. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.016.1>]

- Pérez-Hoyos, A., Martínez, B., García-Haro, F.J., Moreno, A., and Gilabert, M.A., 2014, Identification of ecosystem functional types from coarse resolution imagery using a self-organizing map approach—a case study for Spain: *Remote Sensing*, v. 6, no. 11, p. 11391-11419. [Also available at <http://dx.doi.org/10.3390/rs6111391>.]
- Pons, X., Pesquer, L., Cristóbal, J., and González-Guerrero, O., 2014, Automatic and improved radiometric correction of Landsat imagery using reference values from MODIS surface reflectance images: *International Journal of Applied Earth Observation and Geoinformation*, v. 33, p. 243-254. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.06.002>.]
- Powell, S.J., Jakeman, A., and Croke, B., 2014, Can NDVI response indicate the effective flood extent in macrophyte dominated floodplain wetlands?: *Ecological Indicators*, v. 45, p. 486-493. [Also available at <http://dx.doi.org/10.1016/j.ecolind.2014.05.009>.]
- Proud, S.R., Zhang, Q., Schaaf, C., Fensholt, R., Rasmussen, M.O., Shisanya, C., Mutero, W., Mbow, C., Anyamba, A., Pak, E., and Sandholt, I., 2014, The normalization of surface anisotropy effects present in SEVIRI reflectances by using the MODIS BRDF method: *IEEE Transactions on Geoscience and Remote Sensing*, v. 52, no. 10, p. 6026-6039. [Also available at <http://dx.doi.org/10.1109/TGRS.2013.2294602>.]
- Prush, V.B., and Lohman, R.B., 2014, Time-varying elevation change at the Centralia coal mine in Centralia, Washington (USA), constrained with InSAR, ASTER, and optical imagery: *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*. [Also available at <http://dx.doi.org/10.1109/JSTARS.2014.2348412>.]
- Qiu, B., Fan, Z., Zhong, M., Tang, Z., and Chen, C., 2014, A new approach for crop identification with wavelet variance and JM distance: *Environmental Monitoring and Assessment*, v. 186, no. 11, p. 7929-7940. [Also available at <http://dx.doi.org/10.1007/s10661-014-3977-1>.]
- Qu, B., Ming, J., Kang, S.C., Zhang, G.S., Li, Y.W., Li, C.D., Zhao, S.Y., Ji, Z.M., and Cao, J.J., 2014, The decreasing albedo of the Zhadang glacier on western Nyainqntanglha and the role of light-absorbing impurities: *Atmospheric Chemistry and Physics*, v. 14, no. 20, p. 11117-11128. [Also available at <http://dx.doi.org/10.5194/acp-14-11117-2014>.]
- Quan, J., Chen, Y., Zhan, W., Wang, J., Voogt, J., and Li, J., 2014, A hybrid method combining neighborhood information from satellite data with modeled diurnal temperature cycles over consecutive days: *Remote Sensing of Environment*, v. 155, p. 257-274. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.034>.]
- Reeves, M.C., Moreno, A.L., Bagne, K.E., and Running, S.W., 2014, Estimating climate change effects on net primary production of rangelands in the United States: *Climatic Change*, v. 126, no. 3-4, p. 429-442. [Also available at <http://dx.doi.org/10.1007/s10584-014-1235-8>.]
- Remy, S., and Kaiser, J.W., 2014, Daily global fire radiative power fields estimation from one or two MODIS instruments: *Atmospheric Chemistry and Physics*, v. 14, no. 24, p. 13377-13390. [Also available at <http://dx.doi.org/10.5194/acp-14-13377-2014>.]
- Roupioz, L., Nerry, F., Jia, L., and Menenti, M., 2014, Improved surface reflectance from remote sensing data with sub-pixel topographic information: *Remote Sensing*, v. 6, no. 11, p. 10356-10374. [Also available at <http://dx.doi.org/10.3390/rs61110356>.]
- Roy, A., Royer, A., and Hall, R.J., 2014, Relationship between forest microwave transmissivity and structural parameters for the canadian boreal forest: *IEEE Geoscience and Remote Sensing Letters*, v. 11, no. 10, p. 1802-1806. [Also available at <http://dx.doi.org/10.1109/LGRS.2014.2309941>.]
- Sha, Z., Brown, D.G., Xie, Y., Welsh, W.F., and Bai, Y., 2014, Response of spectral vegetation indices to a stocking rate experiment in Inner Mongolia, China: *Remote Sensing Letters*, v. 5, no. 10, p. 912-921. [Also available at <http://dx.doi.org/10.1080/2150704x.2014.976882>.]
- Shao, Z., and Liu, C., 2014, The integrated use of DMSP-OLS nighttime light and MODIS data for monitoring large-scale impervious surface dynamics—a case study in the Yangtze River Delta: *Remote Sensing*, v. 6, no. 10, p. 9359-9378. [Also available at <http://dx.doi.org/10.3390/Rs6109359>.]

Simms, D.M., Waine, T.W., Taylor, J.C., and Juniper, G.R., 2014, The application of time-series MODIS NDVI profiles for the acquisition of crop information across Afghanistan: International Journal of Remote Sensing, v. 35, no. 16, p. 6234-6254. [Also available at <http://dx.doi.org/10.1080/01431161.2014.951099>.]

Singh, R.K., Senay, G.B., Velpuri, N.M., Bohms, S., and Verdin, J.P., 2014, On the downscaling of actual evapotranspiration maps based on combination of MODIS and Landsat-based actual evapotranspiration estimates: Remote Sensing, v. 6, no. 11, p. 10483-10509. [Also available at <http://dx.doi.org/10.3390/rs61110483>.]

Son, N.T., Chen, C.F., Chen, C.R., Minh, V.Q., and Trung, N.H., 2014, A comparative analysis of multitemporal MODIS EVI and NDVI data for large-scale rice yield estimation: Agricultural and Forest Meteorology, v. 197, p. 52-64. [Also available at <http://dx.doi.org/10.1016/j.agrformet.2014.06.007>.]

Son, Y.-S., Kang, M.-K., and Yoon, W.-J., 2014, Pyrophyllite mapping in the Nohwa deposit, Korea, using ASTER remote sensing data: Geosciences Journal, v. 18, no. 3, p. 295-305. [Also available at <http://dx.doi.org/10.1007/s12303-014-0007-9>.]

Song, X.P., Huang, C., Feng, M., Sexton, J.O., Channan, S., and Townshend, J.R., 2014, Integrating global land cover products for improved forest cover characterization—an application in North America: International Journal of Digital Earth, v. 7, no. 9, p. 709-724. [Also available at <http://dx.doi.org/10.1080/17538947.2013.856959>.]

Srivastava, P.K., Han, D., Rico-Ramirez, M.A., O'Neill, P., Islam, T., and Gupta, M., 2014, Assessment of SMOS soil moisture retrieval parameters using tau-omega algorithms for soil moisture deficit estimation: Journal of Hydrology, v. 519, no. PA, p. 574-587. [Also available at <http://dx.doi.org/10.1016/j.jhydrol.2014.07.056>.]

Startseva, Z., Muzylev, E., Volkova, E., Uspensky, A., and Uspensky, S., 2014, Water and heat regimes modelling for a vast territory using remote-sensing data: International Journal of Remote Sensing, v. 35, no. 15, p. 5775-5799. [Also available at <http://dx.doi.org/10.1080/01431161.2014.945003>.]

Sun, H., Chen, Y., Gong, A., Zhao, X., Zhan, W., and Wang, M., 2014, Estimating mean air temperature using MODIS day and night land surface temperatures: Theoretical and Applied Climatology, v. 118, no. 1-2, p. 81-92. [Also available at <http://dx.doi.org/10.1007/s00704-013-1033-7>.]

Tang, H., Dubayah, R., Brolly, M., Ganguly, S., and Zhang, G., 2014, Large-scale retrieval of leaf area index and vertical foliage profile from the spaceborne waveform lidar (GLAS/ICESat): Remote Sensing of Environment, v. 154, p. 8-18. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.007>.]

Tang, X., Li, H., Desai, A.R., Xu, X., and Xie, J., 2014, Comparison of multiple models for remote sensing of carbon exchange using MODIS data in conifer-dominated forests: International Journal of Remote Sensing, v. 35, no. 24, p. 8252-8271. [Also available at <http://dx.doi.org/10.1080/01431161.2014.981644>.]

Tayebi, M.H., Tangestani, M.H., and Vincent, R.K., 2014, Alteration mineral mapping with ASTER data by integration of coded spectral ratio imaging and SOM neural network model: Turkish Journal of Earth Sciences, v. 23, no. 6, p. 627-644. [Also available at <http://dx.doi.org/10.3906/Yer-1401-9>.]

Teske, R., Giasson, E., and Bagatini, T., 2014, Comparison of the use of digital elevation models in digital mapping of soils of Dois Irmãos, RS, Brazil: Revista Brasileira de Ciencia do Solo, v. 38, no. 5, p. 1367-1376. [Also available at <http://dx.doi.org/10.1590/S0100-06832014000500002>.]

Thackeray, C.W., Fletcher, C.G., and Derksen, C., 2014, The influence of canopy snow parameterizations on snow albedo feedback in boreal forest regions: Journal of Geophysical Research-Atmospheres, v. 119, no. 16, p. 9810-9821. [Also available at <http://dx.doi.org/10.1002/2014jd021858>.]

Thomas, J., Joseph, S., Thrivikramji, K.P., and Arunkumar, K.S., 2014, Sensitivity of digital elevation models—the scenario from two tropical mountain river basins of the Western Ghats, India: Geoscience Frontiers, v. 5, no. 6, p. 893-909. [Also available at <http://dx.doi.org/10.1016/j.gsf.2013.12.008>.]

- Tian, L., Zhang, Y., and Zhu, J., 2014, Decreased surface albedo driven by denser vegetation on the Tibetan Plateau: Environmental Research Letters, v. 9, no. 10. [Also available at <http://dx.doi.org/10.1088/1748-9326/9/10/104001>.]
- Tian, X., Li, Z., Su, Z., Chen, E., van der Tol, C., Li, X., Guo, Y., Li, L., and Ling, F., 2014, Estimating montane forest above-ground biomass in the upper reaches of the Heihe River Basin using Landsat-TM data: International Journal of Remote Sensing, v. 35, no. 21, p. 7339-7362. [Also available at <http://dx.doi.org/10.1080/01431161.2014.967888>.]
- Tomar, V., Mandal, V.P., Srivastava, P., Patairya, S., Singh, K., Ravisankar, N., Subash, N., and Kumar, P., 2014, Rice equivalent crop yield assessment using MODIS Sensors' Based MOD13A1-NDVI Data: IEEE Sensors Journal, v. 14, no. 10, p. 3599-3605. [Also available at <http://dx.doi.org/10.1109/JSEN.2014.2329185>.]
- Tomshin, O.A., and Solovyev, V.S., 2014, The impact of large-scale forest fires on atmospheric aerosol characteristics: International Journal of Remote Sensing, v. 35, no. 15, p. 5742-5749. [Also available at <http://dx.doi.org/10.1080/01431161.2014.945001>.]
- Tripathi, P., Patel, N.R., Kushwaha, S.P.S., and Dadhwal, V.K., 2014, Upscaling of leaf area index in Terai forest plantations using fine- and moderate-resolution satellite data: International Journal of Remote Sensing, v. 35, no. 22, p. 7749-7762. [Also available at <http://dx.doi.org/10.1080/01431161.2014.976886>.]
- Ueyama, M., Ichii, K., Iwata, H., Euskirchen, E.S., Zona, D., Rocha, A.V., Harazono, Y., Iwama, C., Nakai, T., and Oechel, W.C., 2014, Change in surface energy balance in Alaska due to fire and spring warming, based on upscaling eddy covariance measurements: Journal of Geophysical Research-Biogeosciences, v. 119, no. 20, p. 1947-1969. [Also available at <http://dx.doi.org/10.1002/2014jg002717>.]
- Vadrevu, K.P., Lasko, K., Giglio, L., and Justice, C., 2014, Analysis of Southeast Asian pollution episode during June 2013 using satellite remote sensing datasets: Environmental Pollution, v. 195, p. 245-256. [Also available at <http://dx.doi.org/10.1016/j.envpol.2014.06.017>. Epub 2014 Jul 30.]
- Vahmani, P., and Hogue, T.S., 2014, High-resolution land surface modeling utilizing remote sensing parameters and the Noah UCM—a case study in the Los Angeles Basin: Hydrology and Earth System Sciences, v. 18, no. 12, p. 4791-4806. [Also available at <http://dx.doi.org/10.5194/hess-18-4791-2014>.]
- van der Meer, F., Hecker, C., van Ruitenbeek, F., van der Werff, H., de Wijkerslooth, C., and Wechsler, C., 2014, Geologic remote sensing for geothermal exploration—a review: International Journal of Applied Earth Observation and Geoinformation, v. 33, p. 255-269. [Also available at <http://dx.doi.org/10.1016/j.jag.2014.05.007>.]
- Vázquez, P., Calandroni, M., Cabria, F., Agnusdei, M., and Rojas, M.C., 2014, Normalized Difference Vegetation Index (NDVI) patterns in grasslands of the Buenos Aires Province—their association to plant communities and soils: Ecología Austral, v. 24, no. 3, p. 276-285. [Also available at <http://dx.doi.org/http://www.ecologiaustral.com.ar/files/24-3-2.pdf>.]
- Vidot, J., and Borbas, E., 2014, Land surface VIS/NIR BRDF atlas for RTTOV-11—a model and validation against SEVIRI land SAF albedo product: Quarterly Journal of the Royal Meteorological Society, v. 140, no. 684, p. 2186-2196. [Also available at <http://dx.doi.org/10.1002/qj.2288>.]
- Vuolo, F., and Atzberger, C., 2014, Improving land cover maps in areas of disagreement of existing products using NDVI time series of MODIS - Example for Europe: Photogrammetrie, Fernerkundung, Geoinformation, v. 2014, no. 5, p. 393-407. [Also available at <http://dx.doi.org/10.1127/1432-8364/2014/0232>.]
- Wakie, T.T., Evangelista, P.H., Jarnevich, C.S., and Laituri, M., 2014, Mapping current and potential distribution of non-native *prosopis juliflora* in the Afar region of Ethiopia: PLoS ONE, v. 9, no. 11. [Also available at <http://dx.doi.org/10.1371/journal.pone.0112854>.]
- Wang, H., and Ma, M., 2014, Estimation of transpiration and evaporation of different ecosystems in an inland river basin using remote sensing data and the Penman-Monteith equation: Shengtai Xuebao/ Acta Ecologica Sinica, v. 34, no. 19, p. 5617-5626. [Also available at <http://dx.doi.org/10.5846/stxb201301150102>.]

Wang, J., and Ma, Y., 2014, Analysis of Shenyang vegetation change based on remote sensing data of MODIS: Liaoning Gongcheng Jishu Daxue Xuebao (Ziran Kexue Ban)/Journal of Liaoning Technical University (Natural Science Edition), v. 33, no. 11, p. 1484-1487. [Also available at <http://dx.doi.org/10.3969/j.issn.1008-0562.2014.11.010>.]

Wang, L., Zhang, Y., Berninger, F., and Duan, B., 2014, Net primary production of Chinese fir plantation ecosystems and its relationship to climate: Biogeosciences, v. 11, no. 19, p. 5595-5606. [Also available at <http://dx.doi.org/10.5194/bg-11-5595-2014>.]

Wang, P., Gao, F., and Masek, J.G., 2014, Operational data fusion framework for building frequent Landsat-like imagery: IEEE Transactions on Geoscience and Remote Sensing, v. 52, no. 11, p. 7353-7365. [Also available at <http://dx.doi.org/10.1109/TGRS.2014.2311445>.]

Wang, Y., Liu, Y., Li, M., and Tan, L., 2014, The reconstruction of abnormal segments in HJ-1A/B NDVI time series using MODIS—a statistical method: International Journal of Remote Sensing, v. 35, no. 23, p. 7991-8007. [Also available at <http://dx.doi.org/10.1080/01431161.2014.978954>.]

Wang, Y., Shen, Y., Sun, F., and Chen, Y., 2014, Evaluating the vegetation growing season changes in the arid region of northwestern China: Theoretical and Applied Climatology, v. 118, no. 3, p. 569-579. [Also available at <http://dx.doi.org/10.1007/s00704-013-1078-7>.]

Wang, Y., Xie, D., and Li, Y., 2014, Downscaling remotely sensed land surface temperature over urban areas using trend surface of spectral index: Journal of Remote Sensing, v. 18, no. 6, p. 1169-1181. [Also available at <http://dx.doi.org/10.11834/jrs.20144115>.]

Watts, L.M., and Laffan, S.W., 2014, Effectiveness of the BFEST algorithm for detecting vegetation response patterns in a semi-arid region: Remote Sensing of Environment, v. 154, p. 234-245. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.08.023>.]

Weiss, D.J., Atkinson, P.M., Bhatt, S., Mappin, B., Hay, S.I., and Gething, P.W., 2014, An effective approach for gap-filling continental scale remotely sensed time-series: ISPRS Journal of Photogrammetry and Remote Sensing, v. 98, p. 106-118. [Also available at <http://dx.doi.org/10.1016/j.isprsjprs.2014.10.001>.]

Wen, X., Liao, X., Yuan, W., Yan, X., Wei, Z., Liu, H., Feng, J., Lu, S., and Dong, W., 2014, Numerical simulation and data assimilation of the water-energy cycle over semiarid northeastern China: Science China Earth Sciences, v. 57, no. 10, p. 2340-2356. [Also available at <http://dx.doi.org/10.1007/s11430-014-4914-4>.]

Xia, L., Mao, K., Ma, Y., Zhao, F., Jiang, L., Shen, X., and Qin, Z., 2014, An algorithm for retrieving land surface temperatures using VIIRS data in combination with multi-sensors: Sensors (Switzerland), v. 14, no. 11, p. 21385-21408. [Also available at <http://dx.doi.org/10.3390/s141121385>.]

Xiao, J., Ollinger, S.V., Frolking, S., Hurtt, G.C., Hollinger, D.Y., Davis, K.J., Pan, Y., Zhang, X., Deng, F., Chen, J., Baldocchi, D.D., Law, B.E., Arain, M.A., Desai, A.R., Richardson, A.D., Sun, G., Amiro, B., Margolis, H., Gu, L., Scott, R.L., Blanken, P.D., and Suyker, A.E., 2014, Data-driven diagnostics of terrestrial carbon dynamics over North America: Agricultural and Forest Meteorology, v. 197, p. 142-157. [Also available at <http://dx.doi.org/10.1016/j.agrformet.2014.06.013>.]

Xiao, W., Xiong, L., Tang, G., Shao, X., and Wang, J., 2014, Geomorphological divisions of the Tibet Plateau based on topographical feature point groups from DEMs: Annals of GIS, v. 20, no. 4, p. 245-253. [Also available at <http://dx.doi.org/10.1080/19475683.2014.942796>.]

Xie, B., Qin, Z., Wang, Y., and Chang, Q., 2014, Spatial and temporal variation in terrestrial net primary productivity on Chinese Loess Plateau and its influential factors: Nongye Gongcheng Xuebao/Transactions of the Chinese Society of Agricultural Engineering, v. 30, no. 11, p. 244-253. [Also available at <http://dx.doi.org/10.3969/j.issn.1002-6819.2014.11.030>.]

Xu, D., Li, C., Song, X., and Ren, H., 2014, The dynamics of desertification in the farming-pastoral region of North China over the past 10 years and their relationship to climate change and human activity: Catena, v. 123, p. 11-22. [Also available at <http://dx.doi.org/10.1016/j.catena.2014.07.004>.]

- Xu, J., and Shu, H., 2014, Assimilating MODIS-based albedo and snow cover fraction into the Common Land Model to improve snow depth simulation with direct insertion and deterministic ensemble Kalman filter methods: *Journal of Geophysical Research-Atmospheres*, v. 119, no. 18, p. 10684-10701. [Also available at <http://dx.doi.org/10.1002/2014jd022012>.]
- Xu, M., Liang, X.Z., Samel, A., and Gao, W., 2014, MODIS consistent vegetation parameter specifications and their impacts on regional climate simulations: *Journal of Climate*, v. 27, no. 22, p. 8578-8596. [Also available at <http://dx.doi.org/10.1175/Jcli-D-14-00082.1>]
- Xu, Q., Yang, G., Long, H., Wang, C., Li, X., and Huang, D., 2014, Crop information identification based on MODIS NDVI time-series data: *Nongye Gongcheng Xuebao/Transactions of the Chinese Society of Agricultural Engineering*, v. 30, no. 11, p. 134-144. [Also available at <http://dx.doi.org/10.3969/j.issn.1002-6819.2014.11.017>.]
- Xu, Y., Knudby, A., and Ho, H.C., 2014, Estimating daily maximum air temperature from MODIS in British Columbia, Canada: *International Journal of Remote Sensing*, v. 35, no. 24, p. 8108-8121. [Also available at <http://dx.doi.org/10.1080/01431161.2014.978957>.]
- Yan, E.P., Lin, H., Dang, Y.F., and Xia, C.Z., 2014, The spatiotemporal changes of vegetation cover in Beijing-Tianjin sandstorm source control region during 2000—2012: *Shengtai Xuebao/ Acta Ecologica Sinica*, v. 34, no. 17, p. 5007-5020. [Also available at <http://dx.doi.org/10.5846/stxb201305251179>.]
- Yang, L., Cao, Y.G., Zhu, X.H., Zeng, S.H., Yang, G.J., He, J.Y., and Yang, X.C., 2014, Land surface temperature retrieval for arid regions based on Landsat-8 TIRS data—a case study in Shihezi, Northwest China: *Journal of Arid Land*, v. 6, no. 6, p. 704-716. [Also available at <http://dx.doi.org/10.1007/s40333-014-0071-z>.]
- Yao, F., Feng, L., and Zhang, J., 2014, Corn area extraction by the integration of MODIS-EVI time series data and China's environment satellite (HJ-1) data: *Journal of the Indian Society of Remote Sensing*, v. 42, no. 4, p. 859-867. [Also available at <http://dx.doi.org/10.1007/s12524-014-0377-5>.]
- Yu, L., Wang, J., Li, X., Li, C., Zhao, Y., and Gong, P., 2014, A multi-resolution global land cover dataset through multisource data aggregation: *Science China-Earth Sciences*, v. 57, no. 10, p. 2317-2329. [Also available at <http://dx.doi.org/10.1007/s11430-014-4919-z>.]
- Yu, W., Ma, M., Wang, X., Geng, L., Tan, J., and Shi, J., 2014, Evaluation of MODIS LST products using longwave radiation ground measurements in the northern arid region of China: *Remote Sensing*, v. 6, no. 11, p. 11494-11517. [Also available at <http://dx.doi.org/10.3390/rs61111494>.]
- Yuan, W., Cai, W., Liu, S., Dong, W., Chen, J., Arain, M.A., Blanken, P.D., Cescatti, A., Wohlfahrt, G., Georgiadis, T., Genesio, L., Gianelle, D., Grelle, A., Kiely, G., Knohl, A., Liu, D., Marek, M.V., Merbold, L., Montagnani, L., Panferov, O., Peltoniemi, M., Rambal, S., Raschi, A., Varlagin, A., and Xia, J., 2014, Vegetation-specific model parameters are not required for estimating gross primary production: *Ecological Modelling*, v. 292, p. 1-10. [Also available at <http://dx.doi.org/10.1016/j.ecolmodel.2014.08.017>.]
- Yue, C., Ciais, P., Cadule, P., Thonicke, K., Archibald, S., Poulter, B., Hao, W.M., Hantson, S., Mouillet, F., Friedlingstein, P., Maignan, F., and Viovy, N., 2014, Modelling the role of fires in the terrestrial carbon balance by incorporating SPITFIRE into the global vegetation model ORCHIDEE—Part 1—simulating historical global burned area and fire regimes: *Geoscientific Model Development*, v. 7, no. 6, p. 2747-2767. [Also available at <http://dx.doi.org/10.5194/gmd-7-2747-2014>.]
- Zhai, J., Liu, R., Liu, J., Zhao, G., and Huang, L., 2014, Radiative forcing over China due to albedo change caused by land cover change during 1990-2010: *Journal of Geographical Sciences*, v. 24, no. 5, p. 789-801. [Also available at <http://dx.doi.org/10.1007/s11442-014-1120-4>.]
- Zhang, H., Wu, B., Yan, N., Zhu, W., and Feng, X., 2014, An improved satellite-based approach for estimating vapor pressure deficit from MODIS data: *Journal of Geophysical Research-Atmospheres*, v. 119, no. 21, p. 12256-12271. [Also available at <http://dx.doi.org/10.1002/2014JD022118>.]

Zhang, Q., Cheng, Y.-B., Lyapustin, A.I., Wang, Y., Gao, F., Suyker, A., Verma, S., and Middleton, E.M., 2014, Estimation of crop gross primary production (GPP)—fAPAR(chl) versus MOD15A2 FPAR: *Remote Sensing of Environment*, v. 153, p. 1-6. [Also available at <http://dx.doi.org/10.1016/j.rse.2014.07.012>.]

Zhang, X.Q., and Yamaguchi, Y., 2014, Characterization and evaluation of MODIS-derived Drought Severity Index (DSI) for monitoring the 2009/2010 drought over southwestern China: *Natural Hazards*, v. 74, no. 3, p. 2129-2145. [Also available at <http://dx.doi.org/10.1007/s11069-014-1278-1>.]

Zhang, Y., Li, Z., and Li, J., 2014, Comparisons of emissivity observations from satellites and the ground at the CRCs Dunhuang Gobi site: *Journal of Geophysical Research D: Atmospheres*, v. 119, no. 22, p. 13026-13041. [Also available at <http://dx.doi.org/10.1002/2014JD022216>.]

Zhang, Y.-F., Hoar, T.J., Yang, Z.-L., Anderson, J.L., Toure, A.M., and Rodell, M., 2014, Assimilation of MODIS snow cover through the Data Assimilation Research Testbed and the Community Land Model version 4: *Journal of Geophysical Research-Atmospheres*, v. 119, no. 12, p. 7091-7103. [Also available at <http://dx.doi.org/10.1002/2013jd021329>.]

Zhao, J.J., and Liu, L.Y., 2014, Linking satellite-based spring phenology to temperate deciduous broadleaf forest photosynthesis activity: *International Journal of Digital Earth*, v. 7, no. 11, p. 881-896. [Also available at <http://dx.doi.org/10.1080/17538947.2013.786145>.]

Zhao, X., and Liu, Y., 2014, Relative contribution of the topographic influence on the triangle approach for evapotranspiration estimation over mountainous areas, *Advances in Meteorology*, p. 84040-84040.

Zheng, B., Myint, S.W., and Fan, C., 2014, Spatial configuration of anthropogenic land cover impacts on urban warming: *Landscape and Urban Planning*, v. 130, no. 1, p. 104-111. [Also available at <http://dx.doi.org/10.1016/j.landurbplan.2014.07.001>.]

Zhong, B., Zhang, Y., Du, T., Yang, A., Lv, W., and Liu, Q., 2014, Cross-calibration of HJ-1/CCD over a desert site using Landsat ETM + imagery and ASTER GDEM product: *IEEE Transactions on Geoscience and Remote Sensing*, v. 52, no. 11, p. 7247-7263. [Also available at <http://dx.doi.org/10.1109/TGRS.2014.2310233>.]

Zhong, B., Zhang, Y., Du, T., Yang, A., Lv, W., and Liu, Q., 2014, Cross-Calibration of HJ-1/CCD Over a Desert Site Using Landsat ETM plus Imagery and ASTER GDEM Product: *IEEE Transactions on Geoscience and Remote Sensing*, v. 52, no. 11, p. 7247-7263. [Also available at <http://dx.doi.org/10.1109/tgrs.2014.2310233>.]

Zhou, W., Li, J.L., Mu, S.J., Gang, C.C., and Sun, Z.G., Effects of ecological restoration-induced land-use change and improved management on grassland net primary productivity in the Shiyanghe River Basin, north-west China: *Grass and Forage Science*, v. 69, no. 4, p. 596-610. [Also available at <http://dx.doi.org/10.1111/gfs.12073>.]